



Southeast Asian Fisheries
Development Council
Training Department

The 64th Short-term Training Course for University Students

“Ecosystem-Based Fisheries for Sustainable Fisheries Resources Management”




Hokkaido University
Fisheries

Oceanography in the Gulf of Thailand

Hiroji Onishi

13th of May 2024, 13:00~14:00

onishi@fish.hokudai.ac.jp



Gulf of Thailand

Many Tourist Spots



Many Fishing Products

Geographical Feature

Area: 320,000 km²

Length: 800 km

Width: 560 km

Max. Depth: 85 m

Ave. Depth: 58 m


Water Vol.: 12,510 km³

(Stansfield & Garrett 1997)

Gulf of Thailand



South China Sea




Environmental Condition
(Sun Light, Temperature, Air, Water, Flow, Topographies, etc.)

Ecological Pyramid

Land

Sea



Producer
(Primary Production) \equiv

Plants (Photosynthesis)


On Land

In Water

Periphyton

Algae

Phytoplankton



Land Tree

Seed Plants




Land Grass



Seaweed (Giant Kelp)




Seagrass



藍藻類

Blue algae

10µm



鞭毛藻類

Dinoflagellate

10 µm




珪藻類

Diatom


10 µm

Plant on Land



- Holding Land
- Intake Water & Nutrients

Sea alga / Seaweed






Holding sea floor

≠





Appressorium

Why the phytoplankton is so small ?

Phytoplankton




Area-to-Volume ratios

Relative Size			
Radius [L: unit]	1	2	4
Area [L ² : unit ²]	1 ² = 1	2 ² = 4	4 ² = 16
Volume [L ³ : unit ³]	1 ³ = 1	2 ³ = 8	4 ³ = 64
Area / Volume	1/1 = 1	4/8 = 1/2	16/64 = 1/4
Conclusions	<ol style="list-style-type: none"> 1. Most area per unit volume 2. Most drag per unit volume 3. Slowest rate of settling 		Least area per unit volume Least drag per unit volume Fastest rate of settling

To keep staying in surface layer
Large surface area par unit volume

Sink Slower

Sink Faster



Sphere's Projected Area = πr^2

Surface Area = $4 \times \pi r^2$


Volume = $\frac{4}{3} \times \pi r^3$
(≈Mass) when same density

Plant on Land

Tall tree can take a lot of **Sunshine** on the reaves, and take **Nutrients (water)** from its root.





Easy to make **photosynthesis**




Sea alga (Phyto Plankton)

Contradictory advantageous conditions







Plant's Leaf




Phyto-Plankton



Sea alga (Phyto Plankton)

Sunshine(Light) can not penetrate into deep layer, Even in very clear sea water...

Contradictory advantageous conditions



In shallow/costal area,

In deep/offshore area,

River Discharge


Sea bottom is in Photic zone

or

Surface Photic Zone



Nutrient rich bottom water



Nutrients

Poor

Nutrients rich water must be supplied to Photic surface zone

Water Movement

- Vertical Mixing
- Upwelling

Physical Process


Wind Mix. Winter Mix. Tidal Mix. Eddy Mix. etc.

Rich

Sunshine

Rich



Poor



(b) LIGHT ZONES

All month average Chlorophyll

1998-2021, Ave-CHL in All Month




Smaller products
in April & May
than other months,
Because of very strong
water stratification
by very warmer surface
temperature

Sea Surface Temperature(°C)

Averaged Monthly data through the 26 years(1993-2018)

Dec. Jan. Feb.
⇒ Cold water

Apr. May
⇒ Hottest



Water Stratification



Stable



Unstable



Global Warming



Stable

Light Density




Heavy Density

Warm
Low Salinity


Hard
to
Mixing

Cold
High Salinity

Wind and/or Cooling




Unstable



Almost Same Density

Tidal Flow





Easy
to
Mixing

For sustainable primary production (Phytoplankton growth), we must think....

- Sun Energy ⇒Maybe too much in Thailand
 - $\text{CO}_2 + \text{H}_2\text{O} \Rightarrow$... Enough in sea water
 - Nutrients Supply Only slightly controllable by human
- ⇒ Water moving

Photonic Synthesis

- ☐ River discharge (Human effects on Land)
- ☐ Water stratification (Mixing)
- ☐ Upwelling (Vertical moving)
- ☐ Tides/Currents/Eddies (Horizontal moving)




If we fail in control nutrients supply...

If we fail in control nutrients supply...




2024, 2, 27
at Ban Sean Beach



In Japan, there were disease caused by environmental contamination
(4 Big Pollution Diseases)

- Itai-Itai Disease: 1910-
- Minamata Disease: 1956-
- Yokkaichi Asthma : 1961-
- Niigata Minamata Disease:1965-

Water pollution
-> Polluted Plankton
-> Polluted Fish
-> Human Disease



In Japanese dictum,


「過ぎたるは、およばざるが如し」
Sugitaruha, Oyobazaruga, gotoshi



“Too much is as bad as too little.”



**Too clear sea water (poor nutrients) ,
it is difficult to keep
a productive ecosystem**





**Too much nutrients
(overweight condition),
it is difficult to keep
a healthy ecosystem**

In Thai,

นกน้อยทำรังแต่พอตัว



นกน้อยทำรังแต่พอตัว

“Little bird makes small nest”



**Non effect can be seen
 in lower salinity area
 by river discharge
 in September & October,
 Because in costal shallow
 area, there is enough nutrients
 by mixing**

To understand about water distribution in the Gulf of Thailand,
Let's draw **contour lines** of water temperature...




Contour Line:
a line drawn on a map connecting points of **equal height**

On Temperature map \Rightarrow **Equal Temperature**

On Salinity map \Rightarrow **Equal Salinity**


Close together Contour lines \Rightarrow **Front area**




● : Grid Point/Observation Point

27.8 : Numerical Value at the Point (Temperature: °C)

At first,
You must
find
border
points







By Rule of
Thumb




No need to use ruler

Note





Let's draw
counter lines



As a first step,
Draw 1 °C interval
(26, 27, 28 °C)

Next step,
Draw 0.2 °C interval
(x.2, x.4, x.6, x.8 °C)







Moving Speed

At same weight (mass),
Warmer water is bigger volume
(Water level is higher)

Warmer condition =>
Higher Sea Surface



By Global warming
Sea water level is rising...

Molecules: **Low** speed moving
Spans between molecules is **narrow**
High content of molecules
=>
High Density
=>
Heavy Weight/Volume

Molecules: **High** speed moving
Spans between molecules is **wide**
Low content of molecules
=>
Low Density
=>
Light Weight/Volume

Sea Surface Height(m)


Averaged Monthly data through the 26 years(1993-2018)

Nov. Dec. Jan. Feb.


⇒ High Level

Jun. Jul. Aug. Sep.




⇒ Low Level



SW Monsoon
wind







Outward
Ekman
Transport





Inward
Ekman
Transport

NE Monsoon
wind



January


July



Even in both season, Highest point locates at central area in the Gulf.

Why?


5m depth in January




Let's try again
draw
counter lines



5m depth in July



5m depth in January



Let's try again
draw
counter lines



Temperature values
are different.

In the central area,
Warmest point
exists.


In the costal area,
Colder points exist.

In both season,
Warmer & High level
Water exist at the center



We must think about tidal motion effects


5m depth in July



Effect of Tide (Tidal Current)


In the Gulf of Thailand
There is no large-scale current system.
Tidal current is dominant.

Rising tide: inward flow
Ebb tide: outward flow
In all depth





Back-and-Forth
Motion
No Water Mixing

Shear flow makes
Water mixing





Mixed layer
is formed
along the
bottom line





Mixed Water
(Middle range temp, Nutrients content)




In coastal area and shelf edge area, surface warm water and deep nutrient-rich water will be mixed by tidal friction bottom layer.








Good condition to make primary production



Another reason



Mixed Water
(Middle range temp, Nutrients content)



Warmer & High level Water exist at the center





Warmer & High level Water exist at the center

Effect of Eddies

Clockwise Eddies are dominant

Typical Flow Pattern





Upward Air-flow
Lifted-up moisture-air makes cloud & rain


Downward Air-flow
Downward dry-air makes fine weather

Clockwise Eddies and downwelling are dominant in Central area of the Gulf

Another reason



Eddies can carry costal water to off-shore vice versa, off-shore water to costal



Wei Yan, et al., 2018


If strong costal eddies exist for long time...
It makes effect to costal topography...







Gulf of Siam
A.D.1686 (2229)




Gulf of Thailand
Present 2024 (2567)



Healthy ecology cycle...will be destroyed by too warmer surface water, too much nutrients.



Coral bleaching Alert level Over 32°C sea water



**By Global warming
Sea water level is rising ...**



Thailand will melt?





Flooding in Bangkok at High Tide (2021 Nov.)

Bangkok will move?

Road at front of
SEAFDEC/TD
2024/1/25
By Dr. Koki Abe



In 2005, Capital city of MYANMAR, has moved from Yangon to Nay Pyi Taw.

Most of reasons...
Military government...

One of reasons...
High tide disaster of Cyclone



In 2024,
Capital city of Indonesia, Jakarta will move
to East Kalimantan.

A lot of area of Jakarta,
is under 0m height

Jakarta, 2007 Feb.




Summary of my lecture

- In the sea, Phyto Plankton is in contradictory advantageous conditions (Surface rich Sunshine \leftrightarrow Deep rich Nutrients).
- Water movement (Mixing, Upwelling, Tide, Eddy, River Discharge, etc.) must be need to solve this contradictory condition.
- ◆ Too warmer surface condition, Too strong water stratification,
⇒ Stop water mixing, Stop nutrient supply, Stop primary production
⇒ Water level: rise, Flooding risk: higher (NE monsoon season)
- ◆ Too much nutrient supply by river discharge from human effects
⇒ Too heavy bloom, Red tide, Blue-Green Algae, etc.



Thank you very much!



ขอบคุณมากครับ

コップン マーク クラッ

Good Luck!

โชคดีนะ

チョークディーナ

See you again.

พบกันใหม่นะ

ポップガンマイナ

